

TF

Native Bovine Apo Transferrin

Catalog No.	CRA145A CRA145B CRA145C	Quantity:	500 mg 1 g 5 g
Alternate Names:	Serotransferrin, Transferrin, Siderophilin, Beta-1-metal-binding globulin, TF, PRO1557, PRO2086, DKFZp781D0156, Apo Transferrin, ATF.		
Description:	<p>Transferrin is the iron-transport protein of vertebrate serum and donates iron to cells through interaction with a specific membrane receptor, CD71. Transferrin appears to be indispensable for most cells growing in tissue culture.</p> <p>It is referred to frequently as a growth factor because, in analogy to other growth factor-receptor interactions, proliferating cells express high numbers of transferrin receptors, and the binding of transferrin to their receptors is needed for cells to initiate and maintain their DNA synthesis. Apart from its role as an iron transport protein transferrin acts as a cytokine and has functions that may not be related to its iron-carrying capacity</p>		
Gene ID:	280705		
Molecular Weight:	Bovine Apo Transferrin is a glycoprotein of approximately 77 kDa		
Formulation:	The protein (1mg/ml) was lyophilized with no additives.		
Purity:	Greater than 97.0% as determined by Cellulose Acetate.		
Applications:	<p>Bovine Transferrin is a crucial component for the cultivation of mammalian cells <i>in-vitro</i>. Bovine Transferrin is Critical for long-term cells growth <i>in-vitro</i>. Bovine Transferrin is used as detoxificant in media by binding contaminating metal ions. Bovine Transferrin is often used as a nutrient in fermentation media for recombinant protein and biopharmaceutical production. Additional common uses of Bovine Transferrin are Molecular weight, Affinity purification of anti-human transferrin antibodies and also as receptor mediated transfection of molecules such as DNA, into cells.</p>		
Human Virus Test:	<p>FDA approved Plasma from each donor has been tested and found negative for antibody to HIV-1, HIV-2, HCV, HBSAG, HBc, ALT and Syphilis.</p> <p>Viral inactivation by pasteurization (60°C for 10 hours) has been validated using three different test viruses, with removal of 8-14.5 logs of virus documented.</p> <p>The purification process has also been found to remove significant additional quantities of virus</p>		
Physical Appearance:	Sterile Filtered white lyophilized (freeze-dried) powder.		
Storage & Stability:	<p>It is recommended to reconstitute the lyophilized Bovine ATF in sterile 18 MΩ-cm water not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.</p> <p>Lyophilized Bovine Apo Transferrin, although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Bovine Apo Transferrin should be stored at 4°C between 2-7 days and for future use below -18°C.</p> <p>Avoid repeated freeze-thaw cycles.</p>		

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

