cellsciences.com

Goat anti Human IgG, IgM and IgA, Affinity purified Monoclonal Antibody

Catalog No.: PS 103

Quantity: 1.0 mg

Specificity
Antibodies were raised in goat to human IgG, IgM and IgG.
Specific antibodies were absorpted by incubation with Sepharose-bound human immuno-globulins. Specific antibodies were eluted by acidic buffer at pH 2.5 followed by neutralisation and dialysis. After repeated binding with immobilized human immunoglobulins a minimum of 65% protein bound.

No crossreaction with albumin fraction of human serum other then IgG, IgM and IgA.
No crossreaction with albumin fraction of other animal species.

Use
Recommended dilution as secondary antibody in ELISA, 1:1000.
Recommended dilution as secondary antibody in dot-blotting, 1:100.

Presentation
1 ml purified goat antiserum at 1.0 mg/ml protein in 50% glycerol, 0.1 M NaCl, 0.01 M sodium phosphate at pH 7.5.

FOR RESEARCH USE ONLY, NOT FOR DRUG, DIAGNOSTIC OR OTHER USE.

Cell Sciences, Inc.
1504 Providence Highway
Unit 11B
Norwood, MA 02062

Toll Free: 888 769-1246
Phone: 781 769-0610
Fax: 781 769-0542

E-mail: info@cellsciences.com
Web Site: www.cellsciences.com