

Human Kidney Tumor Tissue Lysate

Catalog No.	PX355A	Quantity:	100 µg
	PX355B		0.5 mg

Description: Kidney tumor tissue lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl fluoride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin. Tissue and cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The product was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% β-mercaptoethanol.

Source: Human Kidney tissue - 60 year old male

Buffer: Kidney Tumor lysate is supplied in SDS sample buffer containing 5% β-mercaptoethanol.

Species: Human

Reconstitution: During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For lysates with volumes of 200 µL or less, we recommend gently tapping the vial on a hard surface or briefly centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the container's cap.

Applications: Human kidney tumor tissue lysate is ready to load on SDS -PAGE for Western blotting, 20 µg per lane is recommended for mini gel.

Storage & Stability: Lysate is supplied at a concentration of 2 mg/ml. Store at 2-8°C for continuous use. For extended storage, freeze working aliquots at -70°C. Repeated freezing and thawing is not recommended. Under proper storage conditions the shelf life is half a year from the date of receipt.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.